

Generalized permutative representations of Cuntz algebras

Katsunori Kawamura¹

College of Science and Engineering Ritsumeikan University, 1-1-1 Noji Higashi, Kusatsu, Shiga 525-8577, Japan

Abstract

We introduce representations of the Cuntz algebra $\mathcal{O}_N$ which are parameterized by sequences in the set of unit vectors in $\mathbf{C}^N$. These representations are natural generalizations of permutative representations by Bratteli-Jorgensen and Davidson-Pitts. We show their existence, irreducibility, equivalence, uniqueness of irreducible decomposition and decomposition formulae by using parameters of representations.

1 Introduction

Let $\mathcal{O}_N$ be the Cuntz algebra with canonical generators $s_1, \dots, s_N$ and let $(\mathcal{H}, \pi)$ be a representation of $\mathcal{O}_N$ with a cyclic vector Ω which satisfies

$$\pi(s_1)\Omega = \Omega. \quad (1.1)$$

Then $(\mathcal{H}, \pi)$ is irreducible and unique up to unitary equivalence, and it is equivalent to a permutative representation by [3, 6, 7]. The restriction $(\mathcal{H}, \pi|_{\mathcal{O}_N^{U(1)}})$ on the fixed-point subalgebra $\mathcal{O}_N^{U(1)}$ with respect to the $U(1)$ -gauge action is also irreducible. Especially, $\mathcal{O}_2^{U(1)}$ is isomorphic to the CAR algebra $\mathcal{A}$ with canonical generators $\{a_n : n \geq 1\}$ which satisfy $a_n a_m^* + a_m^* a_n = \delta_{nm} I$ and $a_n a_m + a_m a_n = 0$ for each $n, m \geq 1$. Define an embedding φ of $\mathcal{A}$ into $\mathcal{O}_2$ by

$$\varphi(a_1) \equiv s_1 s_2^*, \quad \varphi(a_n) \equiv \sum_{J \in \{1,2\}^{n-1}} s_J s_1 s_2^* \beta_2(s_J^*) \quad (n \geq 2)$$

where $s_J \equiv s_{j_1} \cdots s_{j_{n-1}}$ for $J = (j_1, \dots, j_{n-1})$ and β_2 is an automorphism of $\mathcal{O}_2$ defined by $\beta_2(s_1) \equiv s_1$ and $\beta_2(s_2) \equiv -s_2$. Then $\varphi(\mathcal{A}) = \mathcal{O}_2^{U(1)}$. Further $(\mathcal{H}, \pi \circ \varphi)$ is equivalent to the Fock representation of $\mathcal{A}$ with the vacuum Ω with respect to the annihilator $\{a_n\}_{n \geq 1}$, that is,

$$(\mathcal{H}, \pi \circ \varphi) \sim \text{Fock}, \quad \pi(\varphi(a_n))\Omega = 0 \quad (n \geq 1).$$

¹e-mail:kawamura@kurims.kyoto-u.ac.jp.

According to such facts, there are many applications for representations of the CAR algebra from representations of $\mathcal{O}_N$ [1, 2, 9]. Further permutative representations distinguish unitary equivalence classes of a certain class of endomorphisms of $\mathcal{O}_N$ effectively by their branching laws [8, 11].

By generalizing (1.1) for a unit vector $z = (z_1, \dots, z_N) \in \mathbf{C}^N$, we define a representation $(\mathcal{H}, \pi)$ with a cyclic vector Ω such that

$$\pi(z_1 s_1 + \dots + z_N s_N)\Omega = \Omega. \quad (1.2)$$

We denote such representation by $GP(z)$. Then $GP(z)$ exists uniquely up to unitary equivalence and irreducible for each z . Further $GP(z)$ and $GP(y)$ are equivalent if and only if $z = y$. The equation (1.2) is equivalent to an eigenequation of the Perron-Frobenius operator for a representation of $\mathcal{O}_N$ associated with a dynamical system [10]. In this way, these representations have many applications and their studies are mathematically interesting.

In this article, we generalize such representations as follows: Define $S(\mathbf{C}^N) \equiv \{z \in \mathbf{C}^N : \|z\| = 1\}$, $S(\mathbf{C}^N)^{\otimes k} \equiv \{z^{(1)} \otimes \dots \otimes z^{(k)} : z^{(i)} \in S(\mathbf{C}^N), i = 1, \dots, k\}$ for $k \geq 1$ and $S(\mathbf{C}^N)^\infty \equiv \{(z^{(n)})_{n \in \mathbf{N}} : z^{(n)} \in S(\mathbf{C}^N)\}$. For $z = (z_i)_{i=1}^N \in \mathbf{C}^N$, define $s(z) \equiv z_1 s_1 + \dots + z_N s_N$.

Definition 1.1 *Let $(\mathcal{H}, \pi)$ be a representation of $\mathcal{O}_N$.*

- (i) *For $z = z^{(1)} \otimes \dots \otimes z^{(k)} \in S(\mathbf{C}^N)^{\otimes k}$, $(\mathcal{H}, \pi)$ is $GP(z)$ if there is a unit cyclic vector $\Omega \in \mathcal{H}$ such that $\pi(s(z))\Omega = \Omega$ and $\{\pi(s(z^{(i)}) \dots s(z^{(k)}))\Omega\}_{i=1}^k$ is an orthonormal family in $\mathcal{H}$ where $s(z) \equiv s(z^{(1)}) \dots s(z^{(k)})$.*
- (ii) *For $z = (z^{(n)})_{n \in \mathbf{N}} \in S(\mathbf{C}^N)^\infty$, $(\mathcal{H}, \pi)$ is $GP(z)$ if there is a unit cyclic vector $\Omega \in \mathcal{H}$ such that $\{\pi(s(z^{(n)})^* \dots s(z^{(1)})^*)\Omega : n \in \mathbf{N}\}$ is an orthonormal family in $\mathcal{H}$.*

For both cases, we call Ω the GP vector of $(\mathcal{H}, \pi)$. We call $(\mathcal{H}, \pi)$ in (i) (resp. (ii)) the GP (=generalized permutative) representation with a cycle (resp. a chain).

In § 3, we show that the case (ii) is equivalent to an induced product representation in [4].

Define $S(\mathbf{C}^N)^{\otimes*} \equiv \coprod_{k \geq 1} S(\mathbf{C}^N)^{\otimes k}$ and $S(\mathbf{C}^N)^\# \equiv S(\mathbf{C}^N)^{\otimes*} \cup S(\mathbf{C}^N)^\infty$.

Theorem 1.2 *For any $z \in S(\mathbf{C}^N)^\#$, there is a representation $(\mathcal{H}, \pi)$ of $\mathcal{O}_N$ which is $GP(z)$. Such representation is unique up to unitary equivalence.*

By Theorem 1.2, the symbol $GP(z)$ makes sense as both a representation and an equivalence class of representations of $\mathcal{O}_N$.

We define three periodicities on $S(\mathbf{C}^N)^\#$ as follows. For $z \in S(\mathbf{C}^N)^{\otimes*}$, z is *periodic* if there are $y \in S(\mathbf{C}^N)^{\otimes l}$ and $p \geq 2$ such that z is equal to the tensor power $y^{\otimes p}$ of y . An element $z = (z^{(n)})_{n \in \mathbf{N}}$ in $S(\mathbf{C}^N)^\infty$ is *eventually periodic* if there are positive integers p, M and a sequence $(c_n)_{n \geq M}$ in $U(1) \equiv \{a \in \mathbf{C} : |a| = 1\}$ such that $z^{(n+p)} = c_n z^{(n)}$ for each $n \geq M$. $z = (z^{(n)})_{n \in \mathbf{N}}$ is *asymptotically periodic* if there is a positive integer p such that $\sum_{n=1}^\infty (1 - |z^{(n)}|z^{(n+p)}|) < \infty$. The eventual periodicity is properly stronger than the asymptotic periodicity.

Theorem 1.3 (i) For $z \in S(\mathbf{C}^N)^{\otimes*}$, $GP(z)$ is irreducible if and only if z is nonperiodic.

(ii) For $z \in S(\mathbf{C}^N)^\infty$, $GP(z)$ is irreducible if and only if z is nonasymptotically periodic.

We define an equivalence relation $\sim$ among $S(\mathbf{C}^N)^\#$. For $z = z^{(1)} \otimes \dots \otimes z^{(k)} \in S(\mathbf{C}^N)^{\otimes k}$ and $y \in S(\mathbf{C}^N)^{\otimes l}$, $z \sim y$ if $l = k$ and $y = z^{(\sigma(1))} \otimes \dots \otimes z^{(\sigma(k))}$ for some $\sigma \in \mathbf{Z}_k$. For $z = (z^{(n)})_{n \in \mathbf{N}}$ and $y = (y^{(n)})_{n \in \mathbf{N}}$ in $S(\mathbf{C}^N)^\infty$, $z \sim y$ if there are nonnegative integers p, q such that $\sum_{n=1}^\infty (1 - |z^{(n+p)}|y^{(n+q)}|) < \infty$. For each $z \in S(\mathbf{C}^N)^{\otimes*}$ and $y \in S(\mathbf{C}^N)^\infty$, we define $z \not\sim y$. For two representations $(\mathcal{H}, \pi)$ and $(\mathcal{H}', \pi')$, $(\mathcal{H}, \pi) \sim (\mathcal{H}', \pi')$ means that $(\mathcal{H}, \pi)$ and $(\mathcal{H}', \pi')$ are unitarily equivalent.

Theorem 1.4 For $z, y \in S(\mathbf{C}^N)^\#$, $GP(z) \sim GP(y)$ if and only if $z \sim y$.

By Theorem 1.2, 1.3 and 1.4, the irreducible decomposition of GP representation makes sense.

Theorem 1.5 (i) If $p \geq 2$ and $z \in S(\mathbf{C}^N)^{\otimes*}$ is nonperiodic, then the following irreducible decomposition holds:

$$GP(z^{\otimes p}) \sim GP(\zeta_1 \cdot z) \oplus \dots \oplus GP(\zeta_p \cdot z).$$

where $\zeta_j \equiv e^{2\pi\sqrt{-1}(j-1)/p}$. This decomposition is unique up to unitary equivalence and multiplicity free.

(ii) If $z \in S(\mathbf{C}^N)^\infty$ is eventually periodic, then there is a nonperiodic element $y \in S(\mathbf{C}^N)^{\otimes*}$ such that the following direct integral decomposition holds:

$$GP(z) \sim \int_{U(1)}^\oplus GP(c \cdot y) d\eta(c) \quad (1.3)$$

where η is the Haar measure of $U(1)$. If there is another $y' \in S(\mathbf{C}^N)^{\otimes*}$ which satisfies (1.3) with respect to z , then there is $c_0 \in U(1)$ such that $y' \sim c_0 \cdot y$.

By Theorem 1.5 (ii), it is understood that the reason why cycles and chains are simultaneously treated in this article.

In § 2, we show Theorem 1.2. In § 3, we show states of the Cuntz algebras associated with GP representations and relations between GP representations with a chain and induced product representations. In § 4.1, we prove a part of Theorem 1.3. In § 4.2, we prove Theorem 1.4. In § 4.3, we prove the remaining part of Theorem 1.3 and Theorem 1.5. In § 5, we explain that any cyclic permutative representation is a GP representation and we introduce examples which are neither permutative representation nor the rotation of permutative representation by the canonical action of $U(N)$. At last, we discuss a remaining problem in § 5.3.

2 Existence, basis and uniqueness

For $N \geq 2$, let $\mathcal{O}_N$ be the *Cuntz algebra* [5], that is, a C^* -algebra which is universally generated by generators $s_1, \dots, s_N$ satisfying $s_i^* s_j = \delta_{ij} I$ for $i, j = 1, \dots, N$ and $s_1 s_1^* + \dots + s_N s_N^* = I$. Let α be the canonical $U(N)$ action on $\mathcal{O}_N$. In this article, any representation means a unital $*$ -representation.

Proposition 2.1 *For each $z \in S(\mathbf{C}^N)^\#$, there is a representation of $\mathcal{O}_N$ which is $GP(z)$.*

Proof. (i) Let $z = z^{(1)} \otimes \dots \otimes z^{(k)} \in S(\mathbf{C}^N)^{\otimes k}$ and $z^{(l)} = (z_1^{(l)}, \dots, z_N^{(l)})$ for $l = 1, \dots, k$. Define $Y_k \equiv \{1, \dots, k\} \times \mathbf{N}$ and a representation π_0 of $\mathcal{O}_N$ on $\mathcal{H} \equiv l_2(Y_k)$ by $\pi_0(s_i) e_{l,m} \equiv e_{l-1, N(m-1)+i}$ for $i = 1, \dots, N$ and $(l, m) \in Y_k$ where $\mathbf{N} \equiv \{1, 2, 3, \dots\}$, $\{e_x\}_{x \in Y_k}$ is the canonical basis of $\mathcal{H}$ and we define $e_{0,m} \equiv e_{k,m}$ for $m \in \mathbf{N}$. Because $e_{l,1} = \pi_0(s_1^{l-1})^* e_{1,1}$ for each $l = 2, \dots, k$, $\{e_x\}_{x \in Y_k} \subset \pi_0(\mathcal{O}_N) e_{1,1}$. This implies that $(\mathcal{H}, \pi_0)$ is cyclic. Choose a sequence $(g(l))_{l=1}^k$ in $U(N)$ which satisfies $g(l)_{j1} = z_j^{(l)}$ for each $l = 1, \dots, k$ and $j = 1, \dots, N$ where $g(l) = (g(l)_{ij})_{i,j=1}^N$. Define a new representation π on $\mathcal{H}$ by $\pi(s_i) e_{l,m} \equiv \pi_0(\alpha_{g(l-1)}(s_i)) e_{l,m}$ for $i = 1, \dots, N$ and $(l, m) \in Y_k$ where we define $g(0) \equiv g(k)$. Since $\alpha_{g(l)}(s_1) = s(z^{(l)})$, we have $\pi(s(z^{(l-1)})) e_{l,1} = e_{l-1,1}$ for $l = 1, \dots, k$ where we define $z^{(0)} \equiv z^{(k)}$. From this, $\pi(s(z)) e_{1,1} = e_{1,1}$. Because $\{e_x\}_{x \in Y_k} \subset \pi(\mathcal{O}_N) e_{1,1}$, $(\mathcal{H}, \pi)$ is cyclic. Therefore $(\mathcal{H}, \pi)$ is $GP(z)$.

(ii) Let $z = (z^{(n)})_{n \in \mathbf{N}} \in S(\mathbf{C}^N)^\infty$, $z^{(n)} = (z_1^{(n)}, \dots, z_N^{(n)}) \in S(\mathbf{C}^N)$ and $Y_\infty \equiv \mathbf{Z} \times \mathbf{N}$. Define a representation π_0 of $\mathcal{O}_N$ on $\mathcal{H} \equiv l_2(Y_\infty)$ by $\pi_0(s_i)e_{n,m} \equiv e_{n-1, N(m-1)+i}$ for $i = 1, \dots, N$ and $(n, m) \in Y_\infty$. Then $\{\pi_0((s_1^*)^n)e_{0,1}\}_{n \in \mathbf{N}} = \{e_{n,1}\}_{n \in \mathbf{N}}$. For convenience, we extend z as $z = (z^{(n)})_{n \in \mathbf{Z}}$ by $z^{(-n)} \equiv (1, 0, \dots, 0) \in S(\mathbf{C}^N)$ for each $n \geq 0$. Choose a sequence $(g(n))_{n \in \mathbf{Z}}$ in $U(N)$ which satisfies

$$g(n)_{j1} = z_j^{(n-1)} \quad (j = 1, \dots, N, n \in \mathbf{Z}). \quad (2.1)$$

Define a new representation π of $\mathcal{O}_N$ on $\mathcal{H}$ by $\pi(s_i)e_{n,m} \equiv \pi_0(\alpha_{g(n)}^*(s_i))e_{n,m}$ for $(n, m) \in Y_\infty$. Because $s(z^{(n-1)}) = \alpha_{g(n)}(s_1)$, $\{\pi(s(z[n]))^*e_{0,1}\}_{n \in \mathbf{N}} = \{e_{n,1}\}_{n \in \mathbf{N}}$ is an orthonormal family in $\mathcal{H}$ where $z[n] \equiv z^{(1)} \otimes \dots \otimes z^{(n)}$. Because $\{e_x\}_{x \in Y_\infty} \subset \pi(\mathcal{O}_N)e_{0,1}$, $(\mathcal{H}, \pi)$ is cyclic. Hence $(\mathcal{H}, \pi)$ is $GP(z)$. ■

In order to show the uniqueness of a GP representation, we construct a basis of the representation space of a given GP representation which is unique up to unitary equivalence.

For $z = z^{(1)} \otimes \dots \otimes z^{(k)} \in S(\mathbf{C}^N)^{\otimes k}$, let $(\mathcal{H}, \pi)$ be $GP(z)$ with the GP vector Ω . Define a unit vector

$$E_n \equiv \pi(s(z^{(n)}) \dots s(z^{(k)}))\Omega \quad (n = 1, \dots, k). \quad (2.2)$$

Then $\pi(s(z^{(n-1)}))E_n = E_{n-1}$ for $n = 2, \dots, k$ and $\pi(s(z^{(k)}))E_1 = E_k$. We see that $\langle E_n | E_{n'} \rangle = \delta_{nn'}$ for $n, n' = 1, \dots, k$. Choose an orthonormal family $\{z^{(n,j)}\}_{j=1}^N$ in $\mathbf{C}^N$ such that $z^{(n,1)} = z^{(n)}$, and define

$$E_{n,j} \equiv \pi(s(z^{(n,j)}))E_{n+1} \quad (j = 1, \dots, N, n = 1, \dots, k)$$

where $E_{k+1} \equiv E_1$. Then $E_{n,1} = E_n$ and $\{E_{n,j} : (n, j) \in \{1, \dots, k\} \times \{1, \dots, N\}\}$ is an orthonormal family in $\mathcal{H}$. Let $\{1, \dots, N\}^0 \equiv \{0\}$ and $\{1, \dots, N\}^*$ be the union of $\{1, \dots, N\}^m$ with respect to every $m = 0, 1, 2, \dots$. Define

$$E_{J,n,j} \equiv \pi(s_J)E_{n,j} \quad (J \in \{1, \dots, N\}^*, n = 1, \dots, k, j = 2, \dots, N)$$

where we define $s_0 \equiv I$ for convenience. Then

$$\mathcal{B}(z) \equiv \{E_n\}_{n=1}^k \cup \{E_{J,n,j} : (J, n, j) \in \{1, \dots, N\}^* \times \{1, \dots, k\} \times \{2, \dots, N\}\}$$

is also an orthonormal family in $\mathcal{H}$. Define a subset $\Lambda(z)$ of $S(\mathbf{C}^N)^{\otimes*}$ by

$$\Lambda(z) \equiv \coprod_{m \geq 0} \{\Lambda_1^{(m)}(z) \sqcup \dots \sqcup \Lambda_k^{(m)}(z)\}$$

where $\Lambda_n^{(0)}(z) \equiv \{z^{(n)} \otimes \cdots \otimes z^{(k)}\}$ for $n = 1, \dots, k$, $\Lambda_1^{(1)}(z) \equiv \{z^{(k,j)}\}_{j=2}^N$, $\Lambda_n^{(1)}(z) \equiv \{z^{(n-1,j)} \otimes z^{(n)} \otimes \cdots \otimes z^{(k)}\}_{j=2}^N$ for $n = 2, \dots, k$, $\Lambda_n^{(m)}(z) \equiv \{\varepsilon_J \otimes x : x \in \Lambda_n^{(1)}(z), J \in \{1, \dots, N\}^{m-1}\}$ for $n = 1, \dots, k$ and $m \geq 2$ where $\{\varepsilon_i\}_{i=1}^N$ is the canonical basis of $\mathbf{C}^N$ and $\varepsilon_J \equiv \varepsilon_{j_1} \otimes \cdots \otimes \varepsilon_{j_m}$ when $J = (j_1, \dots, j_m) \in \{1, \dots, N\}^m$.

Proposition 2.2 *For $z \in S(\mathbf{C}^N)^\#$, let $(\mathcal{H}, \pi)$ be $GP(z)$ with the GP vector Ω . Define a family $\mathcal{C}_z \equiv \{E_x \in \mathcal{H} : x \in \Lambda(z)\}$ of unit vectors in $\mathcal{H}$ by*

$$E_x \equiv \pi(s(x))\Omega \quad (x \in \Lambda(z)). \quad (2.3)$$

Then $\mathcal{C}_z$ is a complete orthonormal basis of $\mathcal{H}$.

Proof. We can verify that $\mathcal{C}_z = \mathcal{B}(z)$. Hence $\mathcal{C}_z$ is an orthonormal family in $\mathcal{H}$. On the other hand, $V_z \equiv \text{Lin}\langle \{E_n\}_{n=1}^k \rangle$ satisfies $\pi(s_J^*)V_z \subset V_z$ for each $J \in \{1, \dots, N\}^*$. Furthermore we see that $\overline{\pi(\mathcal{O}_N)V_z} = \mathcal{H}_0 \equiv \overline{\text{Lin}\langle \{\pi(s_J)\Omega : J \in \{1, \dots, N\}^*\} \rangle}$ where the overline means the closure in $\mathcal{H}$. Because Ω is a cyclic vector, $\mathcal{H} = \mathcal{H}_0$. By definition of $\Lambda(z)$, $\text{Lin}\langle \mathcal{C}_z \rangle$ is dense in $\mathcal{H}_0$. In consequence, the statement holds. $\blacksquare$

Fix $z = (z^{(n)})_{n \in \mathbf{N}} \in S(\mathbf{C}^N)^\infty$. We extend as $z = (z^{(n)})_{n \in \mathbf{Z}}$ by $z^{(-n)} \equiv \varepsilon_1$ for $n \geq 0$. Choose $(g(n))_{n \in \mathbf{Z}}$ which satisfies (2.1) and $g(-n) = I$ for $n \geq 0$. Let $(\mathcal{H}, \pi)$ be $GP(z)$ with the GP vector Ω . Define

$$E_{-n} \equiv \pi(s_1^n)\Omega, \quad E_0 \equiv \Omega, \quad E_n \equiv \pi(s(z^{(1)}) \cdots s(z^{(n)}))^*\Omega \quad (n \geq 1). \quad (2.4)$$

Then we see that $\pi(s(z^{(n)}))E_n = E_{n-1}$ for each $n \in \mathbf{Z}$ and $\{E_n\}_{n \in \mathbf{Z}}$ is an orthonormal family in $\mathcal{H}$. Define

$$E_{n,j} \equiv \pi(s(z^{(n+1,j)}))E_{n+1} \quad ((n,j) \in \mathbf{Z} \times \{1, \dots, N\})$$

where $z^{(n,j)} \equiv (g_{j1}(n), \dots, g_{jN}(n)) \in S(\mathbf{C}^N)$ for $n \in \mathbf{Z}$ and $j = 1, \dots, N$. Then $\{E_{n,j} : (n,j) \in \mathbf{Z} \times \{1, \dots, N\}\}$ is an orthonormal family in $\mathcal{H}$. Especially, $E_{n,1} = E_n$ for each $n \in \mathbf{Z}$. Define

$$E_{J,n,j} \equiv \pi(s_J)E_{n+m,j} \quad (J \in \{1, \dots, N\}^m, n \in \mathbf{Z}, m \in \mathbf{N}, j = 2, \dots, N).$$

By the estimation of values of inner products, we see that $\{E_n\}_{n \in \mathbf{Z}} \cup \{E_{J,n,j} : J \in \{1, \dots, N\}^*, n \in \mathbf{Z}, j = 2, \dots, N\}$ is an orthonormal family in $\mathcal{H}$. For $z \in S(\mathbf{C}^N)^\infty$, define

$$\Lambda(z) \equiv \coprod_{(n,m) \in \mathbf{Z} \times \mathbf{N}} \Lambda_m^{(n)}(z),$$

$\Lambda_1^{(n)}(z) \equiv \{(n, 0)\}$, $\Lambda_2^{(n)}(z) \equiv \{(n, z^{(n+1, j)})\}_{j=2}^N$, $\Lambda_m^{(n)}(z) \equiv \{(n, \varepsilon_J \otimes z^{(n+m-1, j)}) : j = 2, \dots, N, J \in \{1, \dots, N\}^{m-2}\}$ for $m \geq 3$. By a proof of similarity to that of Proposition 2.2, the following holds.

Proposition 2.3 *For $z \in S(\mathbf{C}^N)^\infty$, let $(\mathcal{H}, \pi)$ be $GP(z)$ with the GP vector Ω and let $\{E_n\}_{n \in \mathbf{Z}}$ be as in (2.4). Define*

$$E_{(n, y)} \equiv \pi(s(y))E_{n+m-1} \quad ((n, y) \in \Lambda_m^{(n)}(z), (n, m) \in \mathbf{Z} \times \mathbf{N}).$$

Then $\{E_x : x \in \Lambda(z)\}$ is a complete orthonormal basis of $\mathcal{H}$.

Proof of Theorem 1.2. By Proposition 2.1, the existence is shown. Assume that both $(\mathcal{H}, \pi)$ and $(\mathcal{H}', \pi')$ are $GP(z)$. According to the natural correspondence among bases of $\mathcal{H}$ and $\mathcal{H}'$ in Proposition 2.2 and 2.3, we have a unitary U from $\mathcal{H}$ to $\mathcal{H}'$ such that $U\pi(\cdot)U^* = \pi'(\cdot)$. Hence for $z \in S(\mathbf{C}^N)^\#$, any two GP representations of $\mathcal{O}_N$ by z are equivalent. Therefore the uniqueness is proved. $\blacksquare$

Proposition 2.4 *Assume that $z \in S(\mathbf{C}^N)^{\otimes*}$ and $(\mathcal{H}, \pi)$ is a representation of $\mathcal{O}_N$ with a cyclic vector $\Omega \in \mathcal{H}$ such that $\pi(s(z))\Omega = \Omega$. If z is nonperiodic, then $(\mathcal{H}, \pi)$ is $GP(z)$.*

Proof. Assume that $z = z^{(1)} \otimes \dots \otimes z^{(k)}$ is nonperiodic. Define $y_1 \equiv z$, $y_n \equiv z^{(n)} \otimes \dots \otimes z^{(k)} \otimes z^{(1)} \otimes \dots \otimes z^{(n-1)}$ for $n = 2, \dots, k$ and $E_n \equiv \pi(s(z^{(n)}) \dots s(z^{(k)}))\Omega$ for $n = 1, \dots, k$. Then we have $\langle E_n | E_m \rangle = \langle \pi(s(y_n))E_n | \pi(s(y_m))E_m \rangle = \langle y_n | y_m \rangle \langle E_n | E_m \rangle$. By the nonperiodicity of z , $|\langle y_n | y_m \rangle| < 1$ when $n \neq m$. Hence $|\langle E_n | E_m \rangle| = |\langle y_n | y_m \rangle| \cdot |\langle E_n | E_m \rangle| < |\langle E_n | E_m \rangle|$. This implies that $\langle E_n | E_m \rangle = \delta_{n, m}$ for $n, m = 1, \dots, k$. Hence the statement holds. $\blacksquare$

3 States and induced product representations

In the first, we show the relation between GP representations and states of $\mathcal{O}_N$. Next, we review results in [4]. At last, we show a relation between GP representations with a chain and induced product representations.

For $J \in \{1, \dots, N\}^*$, the *length* $|J|$ of J is defined by $|J| = k$ when $J \in \{1, \dots, N\}^k$. For $z \in S(\mathbf{C}^N)^\#$ and $J = (j_1, \dots, j_m) \in \{1, \dots, N\}^m$, define $z(J) \equiv z_{j_1}^{(1)} \dots z_{j_m}^{(m)}$ and $z(0) \equiv 1$ where we define $z_i^{(kn+l)} \equiv z_i^{(l)}$ for each $n \in \mathbf{N}$, $l = 1, \dots, k$ and $i = 1, \dots, N$ when $z = z^{(1)} \otimes \dots \otimes z^{(k)}$.

Proposition 3.1 For $z \in S(\mathbf{C}^N)^\#$, define a state ω_z of $\mathcal{O}_N$ as follows. When $z = z^{(1)} \otimes \cdots \otimes z^{(k)} \in S(\mathbf{C}^N)^{\otimes k}$,

$$\omega_z(s_J s_K^*) \equiv \overline{z(J)} \cdot z(K) \quad (|J| \equiv |K| \text{ mod } k), \quad \omega_z(s_J s_K^*) \equiv 0 \quad (\text{otherwise})$$

for each $J, K \in \{1, \dots, N\}^*$ with the convention $s_J s_K^* \equiv s_K^*$ when $J = \emptyset$. When $z = (z^{(n)})_{n \in \mathbf{N}} \in S(\mathbf{C}^N)^\infty$,

$$\omega_z(s_J s_K^*) \equiv \delta_{|J|, |K|} \cdot \overline{z(J)} \cdot z(K) \quad (J, K \in \{1, \dots, N\}^*). \quad (3.1)$$

Then the GNS representation by ω_z is $GP(z)$.

Proof. Let $(\mathcal{H}, \pi)$ be $GP(z)$ with the GP vector Ω . Define $\omega'(x) \equiv \langle \Omega | \pi(x) \Omega \rangle$ for $x \in \mathcal{O}_N$. Then we see that $\omega' = \omega_z$. By the uniqueness of GNS representation and the cyclicity of GP representation, the statement holds. $\blacksquare$

Let $\mathcal{H}$ be a Hilbert space in $\mathcal{O}_N$ defined by $\mathcal{H} \equiv \text{Lin}\langle \{s_1, \dots, s_N\} \rangle$ with the inner product $\langle s_i | s_j \rangle \equiv \delta_{ij}$ for $i, j = 1, \dots, N$. For $z = (z^{(i)}) \in S(\mathbf{C}^N)^\infty$, define a map $s_r : \mathcal{H}^{\otimes r} \rightarrow \mathcal{H}^{\otimes(r+1)}$ by $s_r(\xi) \equiv \xi \otimes z^{(r+1)}$. Define the inductive limit $\mathcal{H}_{(0)}^z$ of the inductive system $\{(\mathcal{H}^{\otimes r}, s_r)\}_{r \geq 1}$ of Hilbert spaces and a vector $\Omega_{(0)}^z$ in $\mathcal{H}_{(0)}^z$ by $\Omega_{(0)}^z \equiv \lim_r z^{(1)} \otimes \cdots \otimes z^{(r)}$. Define subalgebras $\mathcal{B}_n \equiv C^*\langle \{s_J s_K^* : J, K \in \{1, \dots, N\}^n\} \rangle$ and $\mathcal{O}_N^0 \equiv \overline{\bigcup_{n=1}^\infty \mathcal{B}_n}$ of $\mathcal{O}_N$, and denote $v_L \equiv s_{l_1} \otimes \cdots \otimes s_{l_n} \in \mathcal{H}^{\otimes n}$ for $L = (l_1, \dots, l_n) \in \{1, \dots, N\}^n$. Remark that $\mathcal{O}_N^0$ equals to the fixed-point subalgebra $\mathcal{O}_N^{U(1)}$ of $\mathcal{O}_N$ by the $U(1)$ -gauge action. Define a representation π_n of $\mathcal{B}_n$ on $\mathcal{H}^{\otimes n}$ by

$$\pi_n(s_J s_K^*) v_L \equiv \delta_{KL} \cdot v_J \quad (J, K, L \in \{1, \dots, N\}^n).$$

Then we have the inductive limit representation $a_{(0)}^z$ of $\mathcal{O}_N^{U(1)}$ on $\mathcal{H}_{(0)}^z$ by $\{(\mathcal{H}^{\otimes n}, \pi_n)\}_{n \geq 1}$. Define a state $F_{(0)}^z \equiv \langle \Omega_{(0)}^z | a_{(0)}^z(\cdot) \Omega_{(0)}^z \rangle$.

Definition 3.2 (Definition 2.9, [4]) $(\mathcal{H}^z, a^z)$ is the induced product representation of $\mathcal{O}_N$ by z if $(\mathcal{H}^z, a^z, \Omega^z)$ is the GNS representation of $\mathcal{O}_N$ of F^z where F^z is a state of $\mathcal{O}_N$ defined by $F_{(0)}^z \circ P$ and P is the canonical conditional expectation from $\mathcal{O}_N$ onto $\mathcal{O}_N^{U(1)}$.

In [4], the induced product representation is parametrized by the twosided infinite sequence in $\mathcal{H}$. We reformulate that by the onesided sequence here.

Theorem 3.3 For $z \in S(\mathbf{C}^N)^\infty$, the following holds.

- (i) $a^z \sim a^y$ if and only if there is $k \geq 0$ such that $\sum_{n=1}^{\infty} (1 - |\langle z^{(n)} | y^{(n+k)} \rangle|) < \infty$ or $\sum_{n=1}^{\infty} (1 - |\langle z^{(n+k)} | y^{(n)} \rangle|) < \infty$.
- (ii) a^z is irreducible if and only if $\sum_{n=1}^{\infty} (1 - |\langle z^{(n)} | z^{(n+k)} \rangle|) = \infty$ for any $k \geq 1$.
- (iii) There is a family $\{a_n^z\}_{n \in \mathbf{Z}}$ of (nonzero) subrepresentations of $a^z|_{\mathcal{O}_N^{U(1)}}$ such that $a^z|_{\mathcal{O}_N^{U(1)}} = \bigoplus_{n \in \mathbf{Z}} a_n^z$.
- (iv) a^z is $GP(z)$.

Proof. (i), (ii) and (iii) are shown in Theorem 3.16, Corollary 3.17 and Proposition 2.12 in [4], respectively. For states F^z in Definition 3.2 and ω_z in (3.1), we can verify that $F^z = \omega_z$. This implies (iv). $\blacksquare$

4 Irreducibility, equivalence and decomposition

4.1 Irreducibility

Let $S(\mathbf{C}^N)_P^{\otimes*}$, $S(\mathbf{C}^N)_{EP}^{\infty}$, $S(\mathbf{C}^N)_{AP}^{\infty}$ be sets of all periodic elements in $S(\mathbf{C}^N)^{\otimes*}$, all eventually periodic elements in $S(\mathbf{C}^N)^{\infty}$, all asymptotically periodic elements in $S(\mathbf{C}^N)^{\infty}$, respectively, and let $S(\mathbf{C}^N)_{NP}^{\otimes*}$, $S(\mathbf{C}^N)_{NEP}^{\infty}$, $S(\mathbf{C}^N)_{NAP}^{\infty}$ be their complements, respectively. Then we see that $S(\mathbf{C}^N)_P^{\otimes*} = \{v^{\otimes k} : v \in S(\mathbf{C}^N)_{NP}^{\otimes*}, k \geq 2\}$.

Lemma 4.1 *For $z \in S(\mathbf{C}^N)_{NP}^{\otimes*}$, let $(\mathcal{H}, \pi)$ be $GP(z)$ with the GP vector Ω and let $\{E_x : x \in \Lambda(z)\}$ be the basis of $\mathcal{H}$ in (2.3). Then the following holds.*

- (i) *If $z \in S(\mathbf{C}^N)^{\otimes k}$ and $b \in \mathbf{N}$ is not a multiple of k , then $\{\pi(s(z)^*)\}^m E_x$ goes to 0 when $m \rightarrow \infty$ for any $x \in \Lambda(z) \cap S(\mathbf{C}^N)^{\otimes b}$.*
- (ii) *If $v \in \mathcal{H}$ satisfies that $\langle v | \Omega \rangle = 0$, then $\lim_{m \rightarrow \infty} \{\pi(s(z)^*)\}^m v = 0$.*

Proof. (i) For sufficient large $m \in \mathbf{N}$, there are $c_m \in \mathbf{C}$ and $j \in \{1, \dots, k\}$ such that $|c_m| \leq 1$ and $\{\pi(s(z)^*)\}^m E_x = c_m \cdot E_j$. By the assumption on b , we see that $j \neq 1$. The nonperiodicity of z implies that $c_m \rightarrow 0$ when $m \rightarrow \infty$.

(ii) Because v is written as $\sum_{x \in \Lambda(z)} a_x E_x$, it is sufficient to consider the case $v = \pi(s(x))\Omega$ for $x \in \Lambda(z)$. Assume that $z \in S(\mathbf{C}^N)^{\otimes k}$ and $v = \pi(s(x))\Omega$ for $x \in S(\mathbf{C}^N)^{\otimes b} \cap \Lambda(z)$. When b is not a multiple of k , the statement holds by (i). Assume that $b = kl$ for $l \geq 1$. By assumption, $x \neq z$. By checking

every element $x \in (\Lambda(z) \setminus \{z\}) \cap S(\mathbf{C}^N)^{\otimes b}$, we can verify that $\pi(s(z)^*)^m v = 0$ when $m \geq l$. In consequence, the assertion holds. $\blacksquare$

Corollary 4.2 *Let $z \in S(\mathbf{C}^N)_{NP}^{\otimes k}$ and let $(\mathcal{H}, \pi)$ be a representation of $\mathcal{O}_N$. If $\Omega, \Omega' \in \mathcal{H}$ are cyclic vectors which satisfy that $\pi(s(z))\Omega = \Omega$ and $\pi(s(z))\Omega' = \Omega'$, then there is $c \in \mathbf{C}$ such that $\Omega = c\Omega'$.*

Proof. There are $y \in \mathcal{H}$ and $c \in \mathbf{C}$ such that $\Omega' = c\Omega + y$ and $\langle \Omega | y \rangle = 0$. By assumption, we see that $\{\pi(s(z)^*)\}^n \Omega' = \Omega'$ for each $n \in \mathbf{N}$. By Lemma 4.1 (ii), $\lim_{n \rightarrow \infty} \{\pi(s(z)^*)\}^n y = 0$. Hence $\Omega' = \lim_{n \rightarrow \infty} \{\pi(s(z)^*)\}^n \Omega' = c\Omega$. $\blacksquare$

Proposition 4.3 (i) *If $z \in S(\mathbf{C}^N)_{NP}^{\otimes*}$, then $GP(z)$ is irreducible.*

(ii) *For $z \in S(\mathbf{C}^N)^\infty$, $GP(z)$ is irreducible if and only if $z \in S(\mathbf{C}^N)_{NAP}^\infty$.*

Proof. (i) Let $(\mathcal{H}, \pi)$ be $GP(z)$ with the GP vector Ω . For a nonzero vector $v_0 \in \mathcal{H}$, it is sufficient to show that $\Omega \in \pi(\mathcal{O}_N)v_0$. Because $\mathcal{H}$ is cyclic, there is $x \in \mathcal{O}_N$ such that $\langle \pi(x)\Omega | v_0 \rangle \neq 0$. Therefore we can assume that $\langle \Omega | v_0 \rangle = 1$ by replacing v and $\pi(x^*)v$ and normalizing it. Hence we can denote $v_0 = \Omega + y$ for $y \in \mathcal{H}$ such that $\langle \Omega | y \rangle = 0$. Assume that $z \in S(\mathbf{C}^N)_{NP}^{\otimes k}$. By Lemma 4.1, we see that $\lim_{n \rightarrow \infty} (\pi(s(z)^*))^n v_0 = \Omega$. Hence $\Omega \in \pi(\mathcal{O}_N)v_0$. Therefore $(\mathcal{H}, \pi)$ is irreducible.

(ii) This holds by Theorem 3.3 (ii) and (iv). $\blacksquare$

The inverse of Proposition 4.3 (i) is shown in § 4.3.

4.2 Equivalence

Lemma 4.4 *Let $z, y \in S(\mathbf{C}^N)^{\otimes*}$ and let $(\mathcal{H}, \pi)$ be a representation of $\mathcal{O}_N$. If $z \not\sim y$ and there are $\Omega, \Omega' \in \mathcal{H}$ which satisfy $\pi(s(z))\Omega = \Omega$ and $\pi(s(y))\Omega' = \Omega'$, then $\langle \Omega | \Omega' \rangle = 0$.*

Proof. Assume that $z \in S(\mathbf{C}^N)^{\otimes k}$ and $y \in S(\mathbf{C}^N)^{\otimes l}$. If $k \neq l$, then $\langle \Omega | \Omega' \rangle = \langle \pi(s(z)^l)\Omega | \pi(s(y)^k)\Omega' \rangle = \langle z^{\otimes l} | y^{\otimes k} \rangle \langle \Omega | \Omega' \rangle$. Because $z \not\sim y$, $|\langle z^{\otimes l} | y^{\otimes k} \rangle| < 1$. Hence $\langle \Omega | \Omega' \rangle = 0$. Assume that $k = l$. If $|\langle z | y \rangle| < 1$, then we see that $\langle \Omega | \Omega' \rangle = 0$. If $|\langle z | y \rangle| = 1$, then there is $c \in U(1)$ such that $y = cz$. Since $z \not\sim y$, $c \neq 1$. By $\pi(s(y))\Omega = c\pi(s(z))\Omega = c\Omega$, $\langle \Omega | \Omega' \rangle = 0$. $\blacksquare$

Proposition 4.5 *If $z, y \in S(\mathbf{C}^N)^{\otimes*}$ or $z, y \in S(\mathbf{C}^N)^\infty$, then $GP(z) \sim GP(y)$ if and only if $z \sim y$.*

Proof. Let $z, y \in S(\mathbf{C}^N)^{\otimes*}$. Assume that $z \sim y$ and $z \in S(\mathbf{C}^N)^{\otimes k}$. Let $(\mathcal{H}, \pi)$ be $GP(z)$ with the GP vector Ω . By assumption, there is $n \in \{1, \dots, k\}$ such that $y = z^{(n)} \otimes \dots \otimes z^{(k)} \otimes z^{(1)} \otimes \dots \otimes z^{(n-1)}$. Then $E_n \equiv \pi(s(z^{(n)} \otimes \dots \otimes z^{(k)}))\Omega$ satisfies $\pi(s(y))E_n = E_n$ and E_n is also a cyclic vector. Hence $(\mathcal{H}, \pi)$ is $GP(y)$. By the second statement in Theorem 1.2, $GP(z) \sim GP(y)$.

Assume by contradiction that $z \not\sim y$ and $GP(z) \sim GP(y)$. By assumption, there is a representation $(\mathcal{H}, \pi)$ of $\mathcal{O}_N$ such that there are unit cyclic vectors $\Omega, \Omega' \in \mathcal{H}$ which satisfy $\pi(s(z))\Omega = \Omega$ and $\pi(s(y))\Omega' = \Omega'$. By the basis of $GP(z)$ in (2.3), there are $a_1, \dots, a_k \in \mathbf{C}$ and $v \in \mathcal{H}$ such that $\Omega' = a_1 E_1 + \dots + a_k E_k + v$ where $E_1 = \Omega$ and $\langle E_n | v \rangle = 0$ for each $n = 1, \dots, k$. By Lemma 4.4 and $z \not\sim y$, we see that $\langle \Omega' | E_n \rangle = 0$ for each $n = 1, \dots, k$. Hence $\Omega \in V^\perp$ for $V \equiv \text{Lin}\langle \{E_n\}_{n=1}^k \rangle$. By definition of $\{E_x\}_{x \in \Lambda(z)}$, there is the smallest $m_0 \in \mathbf{N}$ such that $\Omega' \in (V_{m_0})^\perp$ where $V_{m_0} \equiv \text{Lin}\langle \{E_x : x \in \Lambda_n^{(m)}(z), m \leq m_0, n = 1, \dots, k\} \rangle$. On the other hand, we see that $\Omega' = \pi(s(y))\Omega' \in (V_{m_0+1})^\perp$ by definition of $\{E_x\}_{x \in \Lambda(z)}$. This contradicts the choice of m_0 . Hence $GP(z) \not\sim GP(y)$.

If $z, y \in S(\mathbf{C}^N)^\infty$, the statement holds by Theorem 3.3 (i) and (iv). ■

We consider a relation between chain and cycle here. For a sequence $\zeta \equiv (z^{(l)})_{l=1}^k$ in $S(\mathbf{C}^N)$, define $\zeta^\infty \equiv (z^{(n)})_{n \in \mathbf{N}} \in S(\mathbf{C}^N)^\infty$ by $z^{(nk+l)} \equiv z^{(i)}$ for each $n \geq 0$ and $l = 1, \dots, k$. For $\xi = (y^{(l)})_{l=1}^k$, if $z \equiv z^{(1)} \otimes \dots \otimes z^{(k)} = y^{(1)} \otimes \dots \otimes y^{(k)}$, then $\zeta^\infty \sim \xi^\infty$. We denote ζ^∞ by z^∞ when there is no ambiguity. For $z \in S(\mathbf{C}^N)^{\otimes*}$, the symbol z^∞ makes sense as an equivalence class in $S(\mathbf{C}^N)^\infty$. This notation is convenient to describe decomposition of representations in § 4.3. By definition, the following holds immediately.

Lemma 4.6 (i) *If $z \in S(\mathbf{C}^N)_{EP}^\infty$, then there is $y \in S(\mathbf{C}^N)_{NP}^{\otimes*}$ such that $z \sim y^\infty$.*

(ii) *If $z, y \in S(\mathbf{C}^N)_{NP}^{\otimes*}$ satisfy that $z^\infty \sim y^\infty$, then there is $c \in U(1)$ such that $z \sim cy$.*

In order to prove Theorem 1.4, we show a branching law of the restriction of a representation of $\mathcal{O}_N$ on $\mathcal{O}_N^{U(1)}$.

Lemma 4.7 *Let $(\mathcal{H}, \pi)$ be $GP(z)$ of $\mathcal{O}_N$ for $z = z^{(1)} \otimes \dots \otimes z^{(k)} \in S(\mathbf{C}^N)_{NP}^{\otimes k}$ with the GP vector Ω . Then the following irreducible decomposition holds.*

$$(\mathcal{H}, \pi|_{\mathcal{O}_N^{U(1)}}) = (V_1, \pi|_{\mathcal{O}_N^{U(1)}}) \oplus \dots \oplus (V_k, \pi|_{\mathcal{O}_N^{U(1)}}) \quad (4.1)$$

where V_i is the completion of $V_{i,0} \equiv \pi(\mathcal{O}_N^{U(1)})e_i$ and $e_i \equiv \pi(s(z^{(i)}) \cdots s(z^{(k)}))\Omega$ for $i = 1, \dots, k$.

Proof. Here we denote $\pi(s_i)$ by s_i simply. Define $E_{JK} \equiv s_J s_K^*$ for $J, K \in \{1, \dots, N\}^l$. Then we see that $\langle E_{JK} e_i | E_{J'K'} e_j \rangle = 0$ when $i \neq j$ for each $J, K \in \{1, \dots, N\}^l$ and $J', K' \in \{1, \dots, N\}^{l'}$. Therefore V_i and V_j are orthogonal when $i \neq j$. On the other hand, if $|J| = kn + i$ for $n \geq 0$ and $i = 0, 1, \dots, k-1$, then let $v \equiv z^{(k-i+1)} \otimes \cdots \otimes z^{(k)} \otimes z^{\otimes n}$. Then $s_J \Omega = s_J s(v)^* e_{k-i+1}$. Because $s_J s(v)^* \in \mathcal{O}_N^{U(1)}$, $s_J \Omega \in V_i$. Because $\text{Lin}(\{s_J \Omega : J \in \{1, \dots, N\}^*\})$ is dense in $\mathcal{H}$, (4.1) holds as a $\mathcal{O}_N^{U(1)}$ -module.

Next we show the irreducibility of V_j . If $x \in V_j$, then there is $J' \in \{1, \dots, N\}^{kl}$ such that $c \equiv \langle e_j | s_{J'}^* x \rangle \neq 0$. We replace x by $c^{-1} \cdot s_{J'}^* x$. Then we have a decomposition $x = e_j + y$ where $\langle y | e_j \rangle = 0$. Define $X \equiv s(z^{(j)}) \cdots s(z^{(k)}) s(z^{(1)}) \cdots s(z^{(j-1)})$ and $T_n \equiv X^n (X^*)^n$ for $n \in \mathbf{N}$. Then $T_n \in \mathcal{O}_N^{U(1)}$ and $T_n x \rightarrow e_j$ when $n \rightarrow \infty$ because z is nonperiodic. Hence $e_j \in \overline{\mathcal{O}_N^{U(1)} x} = V_j$. Therefore V_j is an irreducible $\mathcal{O}_N^{U(1)}$ -module for each j . Hence the statement holds. $\blacksquare$

Proof of Theorem 1.4. By Proposition 4.5, it is sufficient to show that for any $y \in S(\mathbf{C}^N)^{\otimes*}$ and $z \in S(\mathbf{C}^N)^\infty$, $GP(y) \not\sim GP(z)$. By Lemma 4.7 and Theorem 3.3 (iii), branching numbers of components of $GP(y)|_{\mathcal{O}_N^{U(1)}}$ and $GP(z)|_{\mathcal{O}_N^{U(1)}}$ are always finite and infinite, respectively. Hence $GP(y)|_{\mathcal{O}_N^{U(1)}}$ and $GP(z)|_{\mathcal{O}_N^{U(1)}}$ are never equivalent. From this, $GP(y)$ and $GP(z)$ are also never equivalent. $\blacksquare$

4.3 Decomposition

Lemma 4.8 *Let $(\mathcal{H}, \pi)$ be a representation of $\mathcal{O}_N$.*

- (i) *Assume that there are $\Omega_1, \dots, \Omega_M \in \mathcal{H}$ and $z_1, \dots, z_M \in S(\mathbf{C}^N)_{NP}^{\otimes*}$ such that $\pi(s(z_i))\Omega_i = \Omega_i$ for each $i = 1, \dots, M$. If $z_i \not\sim z_j$ when $i \neq j$, then there is a subrepresentation $(V, \pi|_V)$ of $(\mathcal{H}, \pi)$ such that $(V, \pi|_V) \sim GP(z_1) \oplus \cdots \oplus GP(z_M)$.*
- (ii) *If there are $z_1, \dots, z_M, y_1, \dots, y_{M'} \in S(\mathbf{C}^N)_{NP}^{\otimes*}$ such that $(\mathcal{H}, \pi) \sim \bigoplus_{j=1}^M GP(z_j)$ and $(\mathcal{H}, \pi) \sim \bigoplus_{l=1}^{M'} GP(y_l)$, then $M = M'$ and there is $\sigma \in \mathfrak{S}_M$ such that $z_{\sigma(i)} \sim y_i$ for each $i = 1, \dots, M$.*

Proof. (i) If $M = 1$, then it holds by definition. Assume that $M \geq 2$. Let $V_i \equiv \overline{\pi(\mathcal{O}_N)\Omega_i}$. Then $(V_i, \pi|_{V_i})$ is $GP(z_i)$. Since z_i is nonperiodic, $GP(z_i)$ is irreducible by Proposition 4.3 (i). By assumption and Proposition 4.5, $\pi|_{V_i} \not\sim \pi|_{V_j}$ when $i \neq j$. Hence a subspace $V \equiv \bigoplus_{i=1}^M V_i$ satisfies the condition in the statement

(ii) By Proposition 4.3 (i), both $GP(z_i)$ and $GP(y_j)$ are irreducible for each $i = 1, \dots, M$ and $j = 1, \dots, M'$. This implies that $M = M'$ and there is $\sigma \in \mathfrak{S}_M$ such that $GP(y_i) \sim GP(z_{\sigma(i)})$ for each $i = 1, \dots, M$. By Proposition 4.5, $z_{\sigma(i)} \sim y_i$ for each $i = 1, \dots, M$. $\blacksquare$

Lemma 4.9 *For $v \in S(\mathbf{C}^N)^{\otimes*}$ and $p \geq 2$, let $(\mathcal{H}, \pi)$ be $GP(v^{\otimes p})$ with the GP vector Ω . Then $W \equiv \text{Lin}\langle\{\pi(s(v^{\otimes j}))\Omega\}_{j=1}^p\rangle$ is spanned by eigenvectors $\Omega_1, \dots, \Omega_p$ of $\pi(s(v))$ with eigenvalues $1, e^{2\pi\sqrt{-1}/p}, \dots, e^{2\pi\sqrt{-1}(p-1)/p}$.*

Proof. By definition, $\dim W = p$ and $A \equiv \pi(s(v))|_W$ is an action of $\mathbf{Z}_p$ on W . Therefore W is decomposed into subspaces $W_1, \dots, W_p$ as the irreducible decomposition. Choose $h_i \in W_i$ such that $h_i \neq 0$. Then there is $\xi_i \in \mathbf{C}$ such that $Ah_i = \xi_i h_i$ for each $i = 1, \dots, p$. A representation W of $\mathbf{Z}_p$ is cyclic if and only if any component in the irreducible decomposition of W has multiplicity 1. Hence $\{\xi_l\}_{l=1}^p = \{e^{2\pi\sqrt{-1}l/p}\}_{l=1}^p$. $\blacksquare$

Proposition 4.10 *If $(\mathcal{H}, \pi)$ is $GP(v^{\otimes p})$ for $p \geq 2$ and $v \in S(\mathbf{C}^N)^{\otimes*}_{NP}$, then*

$$(\mathcal{H}, \pi) \sim GP(v) \oplus GP(e^{2\pi\sqrt{-1}/p}v) \oplus \dots \oplus GP(e^{2\pi\sqrt{-1}(p-1)/p}v).$$

This decomposition is unique up to unitary equivalence.

Proof. By Lemma 4.9, there are normalized eigenvectors $\Omega_1, \dots, \Omega_p$ of $\pi(s(v))$ in $\mathcal{H}$ with eigenvalues $1, e^{2\pi\sqrt{-1}/p}, \dots, e^{2\pi\sqrt{-1}(p-1)/p}$, respectively. Define $z_i \equiv e^{-2\pi\sqrt{-1}(i-1)/p}v$. Then we have $\pi(s(z_i))\Omega_i = \Omega_i$ for $i = 1, \dots, p$. By Lemma 4.8 (i), there is a subrepresentation $(V, \pi|_V)$ of $(\mathcal{H}, \pi)$ such that

$$(V, \pi|_V) \sim GP(v) \oplus GP(e^{2\pi\sqrt{-1}/p}v) \oplus \dots \oplus GP(e^{2\pi\sqrt{-1}(p-1)/p}v). \quad (4.2)$$

On the other hand, the GP vector Ω of $(\mathcal{H}, \pi)$ belongs to $\text{Lin}\langle\{\Omega_i\}_{i=1}^p\rangle \subset V$ by definition of Ω_i in Lemma 4.9. Because Ω is a cyclic vector, $\mathcal{H} = \overline{\pi(\mathcal{O}_N)\Omega} \subset V \subset \mathcal{H}$. From this and (4.2), the first statement holds. The uniqueness holds by Lemma 4.8 (ii). $\blacksquare$

By Proposition 4.3 (i) and Proposition 4.10, the GP representation of $\mathcal{O}_N$ with a cycle is always decomposed into a finite direct sum of irreducible GP representations of $\mathcal{O}_N$ with a cycle.

Proof of Theorem 1.3. (i) By Proposition 4.10, if $z \in S(\mathbf{C}^N)_P^{\otimes*}$, then $GP(z)$ is not irreducible. By Proposition 4.3 (i), the statement holds.
(ii) This holds by Proposition 4.3 (ii). ■

Proof of Theorem 1.5. (i) is shown by Proposition 4.10. We show that for $v \in S(\mathbf{C}^N)^{\otimes*}$,

$$GP(v^\infty) \sim \int_{U(1)}^\oplus GP(cv) d\eta(c) \quad (4.3)$$

where cv denotes the scalar multiple of v by c .

Let $v = v^{(1)} \otimes \cdots \otimes v^{(k)} \in S(\mathbf{C}^N)^{\otimes k}$, $v^{(l)} = (v_1^{(l)}, \dots, v_N^{(l)}) \in S(\mathbf{C}^N)$ for $l = 1, \dots, k$ and $Y_\infty \equiv \mathbf{Z} \times \mathbf{N}$. Choose a sequence $(g(l))_{l=1}^k$ in $U(N)$ which satisfies $g(l)_{j1} = v_j^{(l)}$ for $j = 1, \dots, N$ and $l = 1, \dots, k$. Define a representation π of $\mathcal{O}_N$ on $\mathcal{H} \equiv l_2(Y_\infty)$ by

$$\pi(s_i)e_{kn+l,m} \equiv \sum_{j=1}^N (g(l)^*)_{ji} e_{kn+l-1, N(m-1)+j} \quad ((n, m) \in Y_\infty, l = 1, \dots, k).$$

Because $s(v^{(l)}) = \alpha_{g(l)}(s_1)$, $\pi(s(v^{(l)}))e_{kn+l,1} = e_{kn+l-1,1}$ and $\pi(s(v))e_{kn,1} = e_{k(n-1),1}$ for each $l = 1, \dots, k$ and $n \in \mathbf{Z}$. In the same way of the proof of Proposition 2.1, we see that $(\mathcal{H}, \pi)$ is $GP(v^\infty)$.

Let $Y_k \equiv \{1, \dots, k\} \times \mathbf{N}$ and $\mathcal{K} \equiv l_2(Y_k)$. Define a unitary operator T from $\mathcal{H}$ to $L_2(U(1), \mathcal{K})$ by $Te_{kn+l,m} \equiv \zeta_n \otimes e_{l+1,m}$ for $l = 0, \dots, k-1$ and $(n, m) \in Y_\infty$ where $\zeta_n(c) \equiv c^n$ for $c \in U(1)$. Define a representation $\Pi \equiv \text{Ad}T \circ \pi$ of $\mathcal{O}_N$ on $L_2(U(1), \mathcal{K})$. Then

$$\Pi(s_i)(\zeta_n \otimes e_{l,m}) = \sum_{j=1}^N (g(l-1)^*)_{ji} \cdot \zeta_n \otimes e_{l-1, N(m-1)+j} \quad (l = 2, \dots, k),$$

$$\Pi(s_i)(\zeta_n \otimes e_{1,m}) = \sum_{j=1}^N (g(k)^*)_{ji} \cdot \zeta_{n-1} \otimes e_{k, N(m-1)+j}$$

for $(n, m) \in Y_\infty$ and $i = 1, \dots, N$. For $c \in U(1)$, define a representation π_c of $\mathcal{O}_N$ on $\mathcal{K}$ by

$$\pi_c(s_i)e_{l,m} \equiv \sum_{j=1}^N (g(l-1)^*)_{ji} \cdot e_{l-1, N(m-1)+j} \quad (l = 2, \dots, k),$$

$$\pi_c(s_i)e_{1,m} \equiv \bar{c} \cdot \sum_{j=1}^N (g(k)^*)_{ji} \cdot e_{k, N(m-1)+j}$$

for $i = 1, \dots, N$ and $m \in \mathbf{N}$. Then we can verify that $(\mathcal{K}, \pi_c)$ is $GP(cv)$ with the GP vector $e_{1,1}$. Further $(\Pi(s_i)\phi)(c) = \pi_c(s_i)\phi(c)$ for each $i = 1, \dots, N$,

$\phi \in L_2(U(1), \mathcal{K})$ and $c \in U(1)$. Therefore $GP(v^\infty) \sim \Pi \sim \int_{U(1)}^\oplus \pi_c d\eta(c) \sim \int_{U(1)}^\oplus GP(cv) d\eta(c)$. Hence (4.3) is shown. From this and Lemma 4.6 (i), the first statement of (ii) is proved. Because $GP(v) \circ \gamma_{\bar{c}^{1/k}} = GP(cv)$ for $c \in U(1)$, (4.3) is rewritten as

$$GP(v^\infty) \sim \int_{U(1)}^\oplus GP(v) \circ \gamma_{c^{1/k}} d\eta(c)$$

where γ is the gauge action on $\mathcal{O}_N$. If $z \in S(\mathbf{C}^N)_{EP}^\infty$, then there is $y \in S(\mathbf{C}^N)_{NP}^{\otimes k}$ such that $GP(z) \sim GP(y^\infty)$ by Lemma 4.6 (i) and Theorem 1.4. From this,

$$GP(z) \sim \int_{U(1)}^\oplus GP(y) \circ \gamma_{c^{1/k}} d\eta(c). \quad (4.4)$$

If there are $l \geq 1$ and $y_0 \in S(\mathbf{C}^N)^{\otimes l}$ which satisfy (4.4) with respect to z , then $GP(y_0^\infty) \sim \int_{U(1)}^\oplus GP(y_0) \circ \gamma_{c^{1/l}} d\eta(c) \sim GP(z) \sim GP(y^\infty)$. By Theorem 1.4, $y_0^\infty \sim y^\infty$. By Lemma 4.6 (ii), the last statement of (ii) is proved. ■

5 Examples

5.1 Permutative representations

We show that GP representations are generalizations of permutative representations by [3]. A data $(\mathcal{H}, \pi)$ is a *permutative representation* of $\mathcal{O}_N$ if there is a complete orthonormal basis $\{e_n\}_{n \in \Lambda}$ of $\mathcal{H}$ and a family $f = \{f_i\}_{i=1}^N$ of maps on Λ such that $\pi(s_i)e_n = e_{f_i(n)}$ for each $n \in \Lambda$ and $i = 1, \dots, N$. For $J = (j_l)_{l=1}^k$ and $c \in U(1) \equiv \{a \in \mathbf{C} : |a| = 1\}$, a representation $(\mathcal{H}, \pi)$ of $\mathcal{O}_N$ is $P(J; c)$ if there is a unit cyclic vector $\Omega \in \mathcal{H}$ and $c \in U(1)$ such that $\pi(s_J)\Omega = c\Omega$ and $\{\pi(s_{j_l} \cdots s_{j_k})\Omega\}_{l=1}^k$ is an orthonormal family in $\mathcal{H}$. We simply denote $P(J; 1)$ by $P(J)$. For $J = (j_n)_{n \in \mathbf{N}} \in \{1, \dots, N\}^\infty \equiv \{(i_n)_{n \in \mathbf{N}} : i_n \in \{1, \dots, N\}\}$, a representation $(\mathcal{H}, \pi)$ of $\mathcal{O}_N$ is $P(J)$ if there is a unit cyclic vector $\Omega \in \mathcal{H}$ such that $\{\pi((s_{J(n)})^*)\Omega : n \in \mathbf{N}\}$ is an orthonormal family in $\mathcal{H}$ where $J_{(n)} \equiv (j_1, \dots, j_n)$. $(\mathcal{H}, \pi)$ is a *cycle* (*chain*) if there is $J \in \{1, \dots, N\}^k$ (*resp.* $J \in \{1, \dots, N\}^\infty$) such that $(\mathcal{H}, \pi)$ is $P(J)$. Then $(\mathcal{H}, \pi)$ is a cyclic permutative representation of $\mathcal{O}_N$ if and only if $(\mathcal{H}, \pi)$ is a cycle or a chain. We see that $P(J; c) = GP(\bar{c} \cdot \varepsilon_J)$ for $J \in \{1, \dots, N\}^k$ and $c \in U(1)$, and $P(J) = GP((\varepsilon_{j_n})_{n \in \mathbf{N}})$ for $J = (j_n)_{n \in \mathbf{N}} \in \{1, \dots, N\}^\infty$.

Example 5.1 (i) The *standard representation* $(l_2(\mathbf{N}), \pi_S)$ of $\mathcal{O}_N$ is defined by $\pi_S(s_i)e_n \equiv e_{N(n-1)+i}$ for $n \in \mathbf{N}$ and $i = 1, \dots, N$ where

$\{e_n\}_{n \in \mathbf{N}}$ is the canonical basis of $l_2(\mathbf{N})$ [1]. Then $(l_2(\mathbf{N}), \pi_S)$ is $GP(z)$ with the GP vector e_1 for $z = (1, 0, \dots, 0) \in S(\mathbf{C}^N)$. Since z is non-periodic, $(l_2(\mathbf{N}), \pi_S)$ is irreducible.

- (ii) Define a representation $(l_2(\mathbf{N}), \pi)$ of $\mathcal{O}_3$ by $\pi(s_1)e_1 \equiv e_2, \pi(s_1)e_2 \equiv e_1, \pi(s_2)e_1 \equiv e_3, \pi(s_3)e_1 \equiv e_4, \pi(s_2)e_2 \equiv e_5, \pi(s_3)e_2 \equiv e_6, \pi(s_i)e_n \equiv e_{3(n-1)+i}$ for $i = 1, 2, 3, n \geq 3$. Then e_1 is a cyclic vector of $(l_2(\mathbf{N}), \pi)$ and $\pi(s_1s_1)e_1 = e_1$. Therefore $(l_2(\mathbf{N}), \pi)$ is $GP((1, 0, 0) \otimes (1, 0, 0))$. We have the irreducible decomposition $l_2(\mathbf{N}) = V_1 \oplus V_2$ where $V_1 \equiv \overline{\pi(\mathcal{O}_3)(e_1 + e_2)}$ and $V_2 \equiv \overline{\pi(\mathcal{O}_3)(e_1 - e_2)}$.

5.2 Non permutative representations

Example 5.2 Define a representation $(L_2[0, 1], \pi_B)$ of $\mathcal{O}_N$ by

$$(\pi_B(s_i)\phi)(x) \equiv \chi_{[(i-1)/N, i/N]}(x)\phi(Nx - i + 1)$$

for $\phi \in L_2[0, 1]$, $x \in [0, 1]$ and $i = 1, \dots, N$ where χ_Y is the characteristic function of a subset Y of $[0, 1]$. Then $(L_2[0, 1], \pi_B)$ is $GP(z)$ for $z \equiv (N^{-1/2}, \dots, N^{-1/2})$. $(L_2[0, 1], \pi_B)$ is also irreducible. Further $(L_2[0, 1], \pi_B \circ \gamma_{\bar{c}})$ is $GP(cz)$ for $c \in U(1)$.

Example 5.3 (i) For a unitary matrix $g = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in U(2)$, let $(l_2(\mathbf{N}), \pi)$ be a representation of $\mathcal{O}_2$ defined by

$$\begin{aligned} \pi(s_1)e_1 &\equiv ae_2 + ce_3, & \pi(s_1)e_2 &\equiv e_1, & \pi(s_2)e_1 &\equiv be_2 + ed_3, & \pi(s_2)e_2 &\equiv e_4, \\ \pi(s_1)e_n &\equiv e_{2n-1}, & \pi(s_2)e_n &\equiv e_{2n} & (n \geq 3). \end{aligned}$$

Then $(l_2(\mathbf{N}), \pi)$ satisfies that $\pi(s_1(\alpha s_1 + \beta s_2))e_1 = e_1$ for $g^{-1} = \begin{pmatrix} \alpha & \gamma \\ \beta & \delta \end{pmatrix} \in U(2)$. Hence this representation is $GP(z)$ for $z \equiv \varepsilon_1 \otimes (\alpha \varepsilon_1 + \beta \varepsilon_2)$. If $\alpha\beta \neq 0$, then this representation does not belong to the class of representation by [3, 6, 7]. We see that $\beta \neq 0$ if and only if z is nonperiodic. This is equivalent that $GP(z)$ is irreducible. If $\beta = 0$, then $|\alpha| = 1$ and the following equivalence holds.

$$GP((\alpha^{1/2}\varepsilon_1)^{\otimes 2}) \sim GP(\alpha^{1/2}\varepsilon_1) \oplus GP(-\alpha^{1/2}\varepsilon_1).$$

Note that this representation is determined only by $(\alpha, \beta) \in \mathbf{C}^2$, $|\alpha|^2 + |\beta|^2 = 1$ up to unitary equivalence. Furthermore for any two different elements in

$S(\mathbf{C}^2)$, the associated representations are inequivalent. In this sense, $S(\mathbf{C}^2)$ is a parameter space of unitary equivalence classes of representations of $\mathcal{O}_2$ in this case.

(ii) Because any $z \in S(\mathbf{C}^N)$ is nonperiodic, a cyclic representation of $\mathcal{O}_N$ with the cyclic vector Ω which satisfies $\pi(s(z))\Omega = \Omega$ is irreducible by Proposition 4.3 (i). For $z, y \in S(\mathbf{C}^N)$, $GP(z) \sim GP(y)$ if and only if $z = y$.

(iii) Let $p \geq 1$ and let $(\mathcal{H}, \pi)$ be a representation of $\mathcal{O}_N$ with a cyclic vector Ω . If $(\mathcal{H}, \pi)$ satisfies the following condition, then $(\mathcal{H}, \pi)$ is irreducible:

$$\pi((s_1 + s_2)(s_1 + \xi s_2)(s_1 + \xi^2 s_2) \cdots (s_1 + \xi^{p-1} s_2))\Omega = 2^{p/2}\Omega$$

where $\xi \equiv e^{2\pi\sqrt{-1}/p}$.

Example 5.4 For $\theta \in [0, 1)$, define $z_\theta = (z_\theta^{(n)})_{n \in \mathbf{N}} \in S(\mathbf{C}^2)^\infty$ by

$$z_\theta^{(n)} \equiv (\cos 2\pi n\theta, \sin 2\pi n\theta) \in S^1 \equiv \{(x, y) \in \mathbf{R}^2 : x^2 + y^2 = 1\} \quad (n \in \mathbf{N}).$$

Then $GP(z_\theta)$ is irreducible if and only if θ is irrational. If there is $p_\theta \equiv \min\{q \in \mathbf{N} : q\theta \in \mathbf{N} \cup \{0\}\}$, then

$$GP(z_\theta) \sim \int_{U(1)}^\oplus GP(c \cdot z_\theta^{(1)} \otimes \cdots \otimes z_\theta^{(p_\theta)}) d\eta(c).$$

Proof. Because

$$\sum_{n=1}^M (1 - |\langle z_\theta^{(n)} | z_\theta^{(n+p)} \rangle|) = 2M \cdot \sin^2 \pi p\theta \quad (p, M \geq 1),$$

we see that there is $p \geq 1$ such that $\sum_{n=1}^\infty (1 - |\langle z_\theta^{(n)} | z_\theta^{(n+p)} \rangle|) < \infty$ if and only if $\theta \in \mathbf{Q}$. Therefore z_θ is asymptotically periodic if and only if $\theta \in \mathbf{Q}$. By Theorem 1.3 (ii), the first statement holds. If p_θ exists, then z_θ is eventually periodic. Hence the last statement holds by (4.3). $\blacksquare$

5.3 Eventual periodicity and asymptotic periodicity

In order to show a remaining problem, we introduce an example $z \in S(\mathbf{C}^2)^\infty$ which is asymptotically periodic but not eventually periodic.

Lemma 5.5 Define positive sequences $\{a_n\}$ and $\{b_n\}$ by

$$c_n \equiv \frac{\pi}{4} - Y_n, \quad d_n \equiv \frac{\pi}{4} + Y_n, \quad Y_n \equiv \frac{1}{2} \sin^{-1} \frac{1}{\sqrt{2n}} \quad (n \geq 1)$$

and let $z \equiv (z^{(n)})_{n \geq 1} \in S(\mathbf{C}^2)^\infty$ by

$$z^{(2n-1)} \equiv (\cos c_n, \sin c_n), \quad z^{(2n)} \equiv (\cos d_n, \sin d_n) \quad (n \geq 1).$$

Then the following holds.

- (i) z is noneventually periodic and asymptotically periodic.
- (ii) z is equivalent to $(1/\sqrt{2}, 1/\sqrt{2})^\infty \in S(\mathbf{C}^2)^\infty$.

Proof. (i) Since z is a sequence in $\mathbf{R}^2$ with positive components and $z^{(n)} \neq z^{(m)}$ when $n \neq m$, it is noneventually periodic. We see that $1 - |\langle z^{(2n-1)} | z^{(2n)} \rangle| = 1/n^2$. Because $d_n - c_{n+1} < d_n - c_n$, we can verify that $1 - |\langle z^{(2n)} | z^{(2n+1)} \rangle| < 1/n^2$. Hence $\sum_{n=1}^\infty (1 - |\langle z^{(n)} | z^{(n+1)} \rangle|) < \pi^2/3 < \infty$. Therefore z is asymptotically periodic.

(ii) Let $v \equiv (1/\sqrt{2}, 1/\sqrt{2})$. Then we see that

$$1 - |\langle z^{(2n-1)} | v \rangle| = 2 \sin^2(Y_n/2) < 2 \sin^2(\sin^{-1}(1/\sqrt{2}n)) = 1/n^2.$$

As well as this, $1 - |\langle z^{(2n)} | v \rangle| < 1/n^2$. In consequence, $\sum_{n=1}^\infty (1 - |\langle z^{(n)} | v \rangle|) < \pi^2/3$. Hence the statement holds. $\blacksquare$

By Lemma 5.5 (i), we have the following tabular about chain case:

noneventually periodic		eventually periodic
indecomposable	gray zone	decomposable
nonasymptotically periodic	asymptotically periodic	
irreducible	non irreducible	

where the decomposability is defined as the style of (4.4). In the “gray zone” $= S(\mathbf{C}^N)_{NEP}^\infty \cap S(\mathbf{C}^N)_{AP}^\infty$, it is not known that whether $GP(z)$ is decomposable or not. According to Lemma 5.5 (ii), we propose the following.

Conjecture 5.6 *If z is asymptotically periodic, then there is always $y \in S(\mathbf{C}^N)_{EP}^\infty$ such that $z \sim y$.*

Acknowledgement: The author would like to thank Prof. Akira Asada for a suggestion to study representations of Cuntz algebras. Florin Ambro and Takeshi Nozawa help us by critical reading of this article.

References

- [1] M. Abe and K. Kawamura, *Recursive fermion system in Cuntz algebra. I —Embeddings of fermion algebra into Cuntz algebra —*, Comm. Math. Phys., **228**, 85-101 (2002).
- [2] ———, *Pseudo Cuntz algebra and recursive FP ghost system in string theory*, Int. J. Mod. Phys., **A18**, No. 4 (2003) 607-625.
- [3] O. Bratteli and P. E. T. Jorgensen, *Iterated function systems and permutation representations of the Cuntz algebra*, Memoirs Amer. Math. Soc., **139** (1999), no.663.
- [4] W. R. Bergmann, R. Conti, *Induced product representation of extended Cuntz algebras*, Annali di Matematica **182**, 271-286 (2003).
- [5] J. Cuntz, *Simple C^* -algebras generated by isometries*, Comm. Math. Phys., **57**, 173-185 (1977).
- [6] K. R. Davidson and D. R. Pitts, *The algebraic structure of non-commutative analytic Toeplitz algebras*, Math. Ann., **311**, 275-303 (1998).
- [7] ———, *Invariant subspaces and hyper-reflexivity for free semigroup algebras*, Proc. London Math. Soc., (3) **78** (1999) 401-430.
- [8] K. Kawamura, *Polynomial endomorphisms of the Cuntz algebras arising from permutations. I —General theory—*, Lett. Math. Phys., **71**, 149-158 (2005).
- [9] ———, *Extensions of representations of the CAR algebra to the Cuntz algebra $\mathcal{O}_2$ —the Fock and the infinite wedge—*, J. Math. Phys., **46** (2005) no. 7, 073509, 12 pp.
- [10] ———, *The Perron-Frobenius operators, invariant measures and representations of the Cuntz-Krieger algebras*, J. Math. Phys., **46** (2005), no. 8, 083514, 6 pp.
- [11] ———, *Branching laws for polynomial endomorphisms of Cuntz algebras arising from permutations*, Lett. Math. Phys., to appear.